

Spectrum of cosmic rays with energy above 10^{17} eV

S.P. Knurenko ^a, V.P. Egorova^a, A.V. Glushkov^a, A.A. Ivanov^a, A.D. Krasilnikov^a, I.T. Makarov^a,
A.A. Mikhailov^a, V.V. Olzoyev^a, M.I. Pravdin^a, A.V. Sabourov^a, I.Ye. Sleptsov^a, G.G. Struchkov^a.

^aYu.G.Shafer Institute of Cosmophysical Research and Aeronomy,
31 Lenin Ave., 677980 Yakutsk, Russia

There are some discrepancies in the results on energy spectrum from Yakutsk, AGASA, and HiRes experiments. In this work differential energy spectrum of primary cosmic rays based on the Yakutsk EAS Array data is presented. For the largest events values of S_{600} and axes coordinates have been obtained using revised lateral distribution function. Simulation of converter's response at large distances showed no considerable underestimation of particle density. Complex shape of spectrum in region of $E_0 > 10^{17}$ eV is confirmed. After adjustment of parameters and additional exposition at the Yakutsk array there are three events with energy $E_0 > 10^{20}$ eV.

1. INTRODUCTION

Research into the cosmic rays above 10^{17} eV spectrum shape and into intensity in steep region near 10^{20} eV predicted by Greisen [1], Zatsepin and Kuzmin [2] are of great importance for their sources detection. Results obtained in various experiments [3,4,5] differ from each other by factor 2 and more in absolute intensity, but their shapes are similar. Variations of intensity cannot be described with simple power law. At energies greater than the GZK-cutoff results are inconsistent. At HiRes there is only one event with $E_0 > 10^{20}$ eV and the spectrum is cut-off. AGASA has registered 11 such events ($\theta < 45^\circ$), this could be an evidence for absence of a cutoff.

At the Yakutsk array after recent analysis have been carried out [3] there is only one event with energy estimated to be greater than 10^{20} eV. To explain this contradiction with AGASA A. Watson assumed that at the Yakutsk array such showers are skipped due to inadequate short integration time for large distances from the axis. In this work we have studied affect of particle arrival time distribution at different distances on estimated density for Yakutsk and AGASA. Simulation showed that distortion of estimated density at the Yakutsk array cannot result in such miscalculation. More substantively is the lateral distribution function (LDF) used for axis deter-

mination. We provided axes coordinates determination with adjusted LDF for the largest events, which in average led to increase of S_{600} . Besides, February 18 2004 a new event with energy 10^{20} eV was detected at the Yakutsk array. As a result of these factors there are three events of $E_0 > 10^{20}$ eV registered at the Yakutsk array.

2. DENSITY MEASUREMENT AT LARGE DISTANCES FROM THE AXIS

A shower disk at large distance from the axis ($R > 1500$ m) is quite thick. At Yakutsk array and at AGASA a nearly similar logarithmic RC-converters of the signal from photomultiplier tube (PMT) to digital code with $r \sim 10$ msec are used. At the Yakutsk array for an event to be treated, a coincidence of signals from both detectors within 2 msec is required. Herewith input of converters is closed in 2 msec after coincidence. In the case when shower front is wide, this may result in underestimation of the density. Besides, there is a possibility, that station doesn't operate due to large difference in particles arrival times on different detectors and so some showers might be skipped. These circumstances have been pointed out by Watson in his report [6]. At AGASA input is permanently open and in the case of wide signal this may lead to density overestimation due

to features of converter.

To examine the influence of the effects mentioned above, we have provided simulation of detector's response for distances $R = 1050, 1500$ and 2000 m, based on the particle distribution approximation obtained at AGASA [7]. A coefficient K_R was considered — a ratio between density estimated with RC-convector and the one set with program. For the Yakutsk array, in the case of detectors with large area, registering large particle densities we got following values: $K_{1050} = 1.05$, $K_{1500} = 0.994$, $K_{2000} = 0.76$. Same points for AGASA: $K_{1050} = 1.065$, $K_{1500} = 1.11$, $K_{2000} = 1.2$. At 2000 m distance for Yakutsk there is 25% underestimation, for AGASA — 20% overestimation.

For the real experiment for the shower with $E_0 = 10^{20}$ eV at $R = 2000$ m about 2 particles per detector is expected. Simulation indicated, that in this case underestimation is much less than $K_{2000} = 0.92$. It is connected with the fact that conversion starts only after the first particle hit and at low density the effective thickness of the shower front decreases. Probability of that the station doesn't operate due to gap between operating of two separate detectors is more than 2 mcsec is 8.5% and it is lower by factor 3 than those due to Poisson fluctuations at this density. But considering this circumstance together with steepness of the LDF at large distances we restricted used effective area beyond the bound of array to obtain the intensity of the largest showers.

Simulation showed no significant underestimation of particle density for distances up to 2000 m for density estimating system at the Yakutsk array. In the case of AGASA, when input of RC-convector is constantly open, besides wide distribution, there is an afterpulse contribution to density overestimation from delayed particles (probably neutrons) together with casual additives from background muons. One can conclude from the data in [7] that lagging neutrons can overstate the density by 1.37 already at 500 m and further. Background muons may cause distortions in wide range of axis distances. If one such particle hits within last 10 mcsec of RC-circuit discharge, then resulting density can be overestimated by factor 2

and more independently of the real density. The effects mentioned above are excluded thanks to closing of converter's input in 2 mcsec.

3. ENERGY SPECTRUM SUMMARY

Events selection was provided as described in [3]. Showers with $\theta < 60^\circ$ were used. For determination of the intensity for showers with $E_0 > 4 \cdot 10^{19}$ eV, an extended area together efficient zone outside the array was used.

At standard procedure of axis determination, the Greisen-Linsley approximation of LDF with parameters obtained before at the Yakutsk array is used [8]. It was shown in [9], that for showers with energy greater than 10^{19} eV this LDF badly corresponded with experimental data at $R > 1000$ m from the axis. A modified approximation was proposed:

$$f(r) \sim \left(\frac{R}{R_0}\right)^{-1.3} \cdot \left(1 + \frac{R}{R_0}\right)^{-(b-1.3)} \times \left(1 + \frac{R}{2000}\right)^{-3.5} \quad (1)$$

For showers with $E_0 > 2 \cdot 10^{19}$ eV parameter b does not depend on energy but depends on θ . In this work we provided axes coordinates determination with this adjusted LDF. As a result, in average estimated S_{600} values increased: 10% — for showers with axes lying within array area and 20% on border.

For energy estimation we used adjusted formulas of S_{300} and S_{600} [10]. For upright showers ($\theta = 0^\circ$, depth $X_0 = 1020 \text{ g} \cdot \text{cm}^{-2}$), the following relationships were obtained using calorimetric method:

$$E_0 = (5.66 \pm 1.4) \cdot 10^{17} \times \left[\frac{S_{300}(0^\circ)}{10}\right]^{0.94 \pm 0.02} \text{ eV}, \quad (2)$$

$$E_0 = (4.6 \pm 1.2) \cdot 10^{17} \times [S_{600}(0^\circ)]^{0.98 \pm 0.02} \text{ eV}. \quad (3)$$

Zenith-angular dependency was accepted in form of:

$$S(\theta) = S(0^\circ) \cdot \left[(1 - \beta) \cdot \exp\left(\frac{X_0 - X}{\lambda_e}\right) + \beta \cdot \exp\left(\frac{X_0 - X}{\lambda_m}\right) \right], \quad (4)$$

where $\lambda_e = 200 \text{ g/cm}^2$ — the attenuation length for the soft component (electrons), $\lambda_m = 1000 \text{ g/cm}^2$ — the attenuation length for the hard component (associated with muons), β — contribution from the hard component to full response $S_{300}(0^\circ)$ or $S_{600}(0^\circ)$ at $1020 \text{ g} \cdot \text{cm}^{-2}$ depth.

The following dependencies of parameters β on S_{300} and S_{600} have been carried out from experimental data:

$$\beta_{300} = (0.368 \pm 0.021) \times \left[\frac{S_{300}(0^\circ)}{10} \right]^{-0.185 \pm 0.02}, \quad (5)$$

$$\beta_{600} = (0.62 \pm 0.06) \times [S_{600}(0^\circ)]^{-0.076 \pm 0.03}. \quad (6)$$




Figure 1. Differential energy spectra according to the Yakutsk EAS array, AGASA [4] and HiRes [5].

On Fig.1 the differential energy spectra obtained at the Yakutsk array, AGASA and HiRes

are presented. Results obtained in different experiments correspond quite well in shape but differ in intensity. The data from the Yakutsk array near 10^{19} eV are higher by factor ~ 2.5 than HiRes data and $\sim 30\%$ than AGASA's. This rather is connected with the difference in estimation of the showers energy. For estimation of energy in upright showers at AGASA the dependency between S_{600} and E_0 based upon the model calculations [7] is used. Our formulas (2) and (3) for upright showers estimate the energy (30–40)% higher at $E_0 \approx 5 \cdot 10^{17} \text{ eV}$ and (15–20)% higher at $E_0 \approx 10^{19} \text{ eV}$ than it follows from similar models.

In the region of ultra-high energies the results from Yakutsk and AGASA approach. There are 4 events registered in Yakutsk with adjusted estimated energy exceeding $10^{19.9} \text{ eV}$ (see Table 1). Relative errors in energy estimation resulting from uncertainty of the parameters in formulas and errors in determination of axis coordinates and zenith angle in these individual showers amount from 32% to 46%. If the energy is reduced by one standard error then it slightly exceeds the 10^{20} eV threshold only in one event. Therefore the relic cutoff of the spectrum cannot be rejected based on Yakutsk EAS data.

Similar experimental errors are observed at AGASA. According to [10] their averaged value is about 20%. Taking into account this circumstance a conclusion was made in [11] that yet there are too few events recorded to approve the spectrum cutoff absence. Besides, estimations of the energy at AGASA depend on model conclusions. The affects of observed densities mentioned above are also not considered yet.

The HiRes results are consistent with the GZK-cutoff of the spectrum, the AGASA and Yakutsk data are inconsistent. But because of small statistics and errors in energy estimation while it is impossible to final conclude about this problem. To solve this and investigate of affinities of the particles with energies above GZK-cutoff, data with high statistics and good accuracy in the energy estimation a necessary available.

Table 1

The most energetic events detected with the Yakutsk array (sorted by energy)

N	Date	Time, UT	θ°	$\log E_0$	δE_0 (%)	b°	l°
1	02.18.04	22:20:38	47.7	20.16	42	16.3	140.2
2	05.07.89	22:03:00	58.7	20.14	46	2.7	161.6
3	12.21.77	18:45:00	46.0	20.01	40	50.0	220.6
4	02.15.78	03:35:00	9.6	19.99	32	15.5	102.0

Acknowledgements

This work is supported by INTAS grant №03–51–5113 and Federal Agency of Science and Innovations grant №748.2003.2.

REFERENCES

1. Greisen K. // Phys. Rev. Lett. 1966. V.16. P.748.
2. Zatsepin G.T. and Kuzmin V.A., JETP lett. 4 (1996) 144.
3. Glushkov A.V. et al. // Proc. 28th ICRC, Tsukuba.2003. V.1 P. 389.
4. Sakaki N. et al. // Proc. 27th ICRC. Gamburg. 2001. V.1 P.333.
5. Abu-Zayyad T. et al. // preprint. 2002. astro-ph//0208301.
6. Watson A.A. // Proc. 28th ICRC, Tsukuba. 2003. V.1 P.373.
7. Takeda M. // Astropart. Phys. 2003. V.19 P.477.
8. Efimov N.N. et al. // Catalogue of Highest Energy Cosmic Ray. №3. World Data Center C2, Japan. 1988. 56p.
9. Glushkov A.V. et al., Russ. Nucl. Phys. 63 (2000) 1477.
10. Egorova V.P. et al. // astro-ph/ 0408493
11. Marco D., Blasi P., Olinto A.V. // Astropart. Phys. 2003. V.20. P53.